

# AHRQ Patient Safety Indicators – Areal-level for California, 2005

Office of Statewide Health Planning and Development (OSHPD)

County	Iatrogenic Pneumothorax, Secondary Diagnosis Field	Selected Infections Due to Medical Care, Secondary Diagnosis Field	Postoperative Wound Dehiscence	Postoperative Hemorrhage or Hematoma
	Rates* (# Cases)	Rates (# Cases)	Rates (# Cases)	Rates (# Cases)
<b>Statewide</b>	<b>6.5 (1,720)</b>	<b>23.2 (6,175)</b>	<b>1.7 (442)</b>	<b>14.3 (3,813)</b>
Alameda	7.7 (84)	23.4 (257)	1.6 (17)	14.6 (160)
Alpine				
Amador	6.3 (2)	34.7 (11)		18.9 (6)
Butte	10.2 (17)	31.1 (52)	3.0 (5)	19.7 (33)
Calaveras	10.6 (4)	18.5 (7)	5.3 (2)	34.4 (13)
Colusa		40.5 (6)		6.7 (1)
Contra Costa	9.2 (70)	22.3 (170)	2.4 (18)	16.6 (126)
Del Norte		22.2 (5)		
El Dorado	7.3 (10)	22.7 (31)	1.5 (2)	19.1 (26)
Fresno	8.2 (50)	23.1 (141)	1.6 (10)	14.2 (87)
Glenn	15.1 (3)	20.2 (4)	5.0 (1)	10.1 (2)
Humboldt	9.9 (10)	20.7 (21)	1.0 (1)	18.8 (19)
Imperial	10.1 (11)	19.2 (21)	1.8 (2)	11.0 (12)

# AHRQ Patient Safety Indicators – Areal-level for California, 2005

Office of Statewide Health Planning and Development (OSHPD)

County	Iatrogenic Pneumothorax, Secondary Diagnosis Field  Rates* (# Cases)	Selected Infections Due to Medical Care, Secondary Diagnosis Field  Rates (# Cases)	Postoperative Wound Dehiscence  Rates (# Cases)	Postoperative Hemorrhage or Hematoma  Rates (# Cases)
<b>Statewide</b>	<b>6.5 (1,720)</b>	<b>23.2 (6,175)</b>	<b>1.7 (442)</b>	<b>14.3 (3,813)</b>
Inyo	7.1 (1)			14.2 (2)
Kern	8.5 (45)	29.1 (154)	1.9 (10)	16.3 (86)
Kings	4.8 (5)	11.6 (12)		15.4 (16)
Lake	15.8 (8)	27.7 (14)	2.0 (1)	39.5 (20)
Lassen		10.7 (3)	7.1 (2)	10.7 (3)
Los Angeles	4.7 (340)	25.6 (1,853)	1.5 (107)	12.6 (910)
Madera	12.6 (13)	19.5 (20)	1.0 (1)	9.7 (10)
Marin	5.1 (10)	20.2 (40)	1.0 (2)	16.2 (32)
Mariposa	6.8 (1)	27.2 (4)		6.8 (1)
Mendocino	7.4 (5)	16.2 (11)		17.7 (12)
Merced	9.7 (16)	27.3 (45)	1.8 (3)	14.0 (23)
Modoc	13.3 (1)	13.3 (1)		13.3 (1)
Mono				

# AHRQ Patient Safety Indicators – Areal-level for California, 2005

Office of Statewide Health Planning and Development (OSHPD)

County	Iatrogenic Pneumothorax, Secondary Diagnosis Field  Rates* (# Cases)	Selected Infections Due to Medical Care, Secondary Diagnosis Field  Rates (# Cases)	Postoperative Wound Dehiscence  Rates (# Cases)	Postoperative Hemorrhage or Hematoma  Rates (# Cases)
<b>Statewide</b>	<b>6.5 (1,720)</b>	<b>23.2 (6,175)</b>	<b>1.7 (442)</b>	<b>14.3 (3,813)</b>
Monterey	6.1 (18)	17.5 (52)	2.0 (6)	15.8 (47)
Napa	9.8 (10)	26.4 (27)	1.0 (1)	33.3 (34)
Nevada	16.4 (13)	26.6 (21)	1.3 (1)	16.4 (13)
Orange	5.2 (115)	21.5 (474)	1.3 (29)	11.4 (250)
Placer	8.2 (20)	15.1 (37)	1.6 (4)	9.8 (24)
Plumas	17.3 (3)	5.8 (1)		11.5 (2)
Riverside	5.6 (79)	21.7 (304)	2.1 (29)	12.7 (178)
Sacramento	7.1 (71)	22.5 (225)	1.1 (11)	16.5 (165)
San Benito	7.7 (3)	28.2 (11)		23.0 (9)
San Bernardino	7.7 (106)	25.3 (347)	2.6 (35)	12.0 (165)
San Diego	6.8 (149)	24.2 (530)	1.5 (32)	14.8 (323)
San Francisco	6.8 (43)	19.0 (120)	1.7 (11)	16.8 (106)
San Joaquin	7.6 (36)	16.8 (79)	1.5 (7)	21.9 (103)

# AHRQ Patient Safety Indicators – Areal-level for California, 2005

Office of Statewide Health Planning and Development (OSHPD)

County	Iatrogenic Pneumothorax, Secondary Diagnosis Field	Selected Infections Due to Medical Care, Secondary Diagnosis Field	Postoperative Wound Dehiscence	Postoperative Hemorrhage or Hematoma
	Rates* (# Cases)	Rates (# Cases)	Rates (# Cases)	Rates (# Cases)
<b>Statewide</b>	<b>6.5 (1,720)</b>	<b>23.2 (6,175)</b>	<b>1.7 (442)</b>	<b>14.3 (3,813)</b>
San Luis Obispo	3.9 (8)	15.6 (32)	1.5 (3)	12.2 (25)
San Mateo	2.8 (15)	16.9 (91)	1.3 (7)	11.3 (61)
Santa Barbara	4.9 (15)	13.5 (41)	1.3 (4)	10.9 (33)
Santa Clara	7.0 (89)	16.8 (215)	1.3 (17)	11.3 (145)
Santa Cruz	8.3 (16)	17.1 (33)	1.6 (3)	7.3 (14)
Shasta	10.2 (14)	44.3 (61)	2.9 (4)	21.8 (30)
Sierra		36.3 (1)		
Siskiyou	2.8 (1)	22.4 (8)		22.4 (8)
Solano	8.0 (24)	24.6 (74)	0.7 (2)	19.3 (58)
Sonoma	10.0 (36)	23.7 (85)	2.0 (7)	13.7 (49)
Stanislaus	6.1 (22)	19.3 (69)	2.5 (9)	22.3 (80)
Sutter	3.1 (2)	13.8 (9)		18.4 (12)
Tehama	15.2 (7)	17.4 (8)		13.1 (6)

**AHRQ Patient Safety Indicators –  
Areal-level for California, 2005**

Office of Statewide Health Planning and Development (OSHPD)

County	Iatrogenic Pneumothorax, Secondary Diagnosis Field	Selected Infections Due to Medical Care, Secondary Diagnosis Field	Postoperative Wound Dehiscence	Postoperative Hemorrhage or Hematoma
	Rates* (# Cases)	Rates (# Cases)	Rates (# Cases)	Rates (# Cases)
<b>Statewide</b>	<b>6.5 (1,720)</b>	<b>23.2 (6,175)</b>	<b>1.7 (442)</b>	<b>14.3 (3,813)</b>
Trinity	8.9 (1)	17.7 (2)	8.9 (1)	17.7 (2)
Tulare	3.9 (11)	23.3 (65)	1.8 (5)	19.4 (54)
Tuolumne	2.1 (1)	15.0 (7)	4.3 (2)	30.0 (14)
Ventura	6.9 (40)	20.5 (119)	2.2 (13)	13.9 (81)
Yolo	6.4 (9)	9.9 (14)	2.1 (3)	17.0 (24)
Yuba	6.3 (3)	21.2 (10)	4.2 (2)	21.2 (10)
Unknown/Missing	(34)	(120)	(10)	(87)